

Differential Effects of Falcarinol and Related Aliphatic C₁₇-Polyacetylenes on Intestinal Cell Proliferation

STIG PURUP,^{*,†} ERIC LARSEN,[†] AND LARS P. CHRISTENSEN[‡]

[†]Department of Animal Health, Welfare and Nutrition, Faculty of Agricultural Sciences, Aarhus University, Blichers Allé 20, P.O. Box 50, DK-8830 Tjele, Denmark, and [‡]Institute of Chemical Engineering, Biotechnology and Environmental Technology, Faculty of Engineering, University of Southern Denmark, Niels Bohrs Allé 1, DK-5230 Odense M, Denmark

Quantitative major polyacetylenes of carrots (falcarinol and falcarindiol) and American ginseng roots (falcarinol and panaxydol) were isolated and tested in human intestinal epithelial cells of normal (FHs 74 Int.) and cancer (Caco-2) origin. A hormesis effect was seen for all isolated polyacetylenes when added to Caco-2 cells in concentrations ranging from 1 ng/mL to 20 μ g/mL. The relative inhibitory potency was falcarinol > panaxydol > falcarindiol. No hormesis effect was observed when adding the polyacetylenes to FHs 74 Int. cells. Instead, an inhibitory growth response was observed above 1 μ g/mL. The relative inhibitory potency was panaxydol > falcarinol > falcarindiol. Maximal inhibition at 20 μ g/mL corresponded to approximately 95% and 80% inhibition of cell proliferation in normal and cancer cells, respectively. Combinations of falcarinol and falcarindiol added to normal and cancer cells showed a synergistic response for the inhibition of cell growth. Furthermore, the oxidized form of falcarinol, falcarinon, showed a significantly less growth inhibitory effect in intestinal cells of both normal and cancer origin; hence, a hydroxyl group at C-3 may be important for activity of falcarinol-type polyacetylenes. Extracts of carrots, containing different amounts of falcarinol, falcarindiol, and falcarindiol 3-acetate had significant inhibitory effects on both normal and cancer cell proliferation. In cancer cells, the extract containing the highest concentration of falcarinol tended to have the highest growth inhibitory effect, in accordance with a higher potency of falcarinol than falcarindiol. The present study demonstrates that aliphatic C₁₇-polyacetylenes are potential anti-cancer principles of carrots and related vegetables and that synergistic interaction between bioactive polyacetylenes may be important for their bioactivity.

KEYWORDS: *Daucus carota*; *Panax quinquefolium*; falcarinol; falcarindiol; panaxydol; falcarinon; polyacetylenes; bioactivity; intestinal epithelial cells; proliferation; synergistic interactions

INTRODUCTION

Epidemiological studies have provided evidence that a diet high in fruit and vegetables is associated with a reduced risk for the development of certain types of cancer, cardiovascular diseases, diabetes, and other diseases (1–7). Compounds associated with the health promoting effects of vegetables are glucosinolates and other organosulfur compounds and their degradation products, carotenoids, phytosterols, polyphenols, vitamins, and dietary fibers (1, 7). The mechanisms for the protection of these classes of natural products are mostly unknown and may only in part explain the health effects of vegetables. Consequently, in recent years focus has been on other types of potential health promoting compounds. One of these groups of compounds is aliphatic C₁₇-polyacetylenes of the falcarinol-type (Figure 1) (7–13), which are common in carrots and related vegetables such as parsley, celery, parsnip, and fennel (10, 11, 14) as well as in medicinal plants such as ginseng (15, 16). Polyacetylenes of the falcarinol-type have shown many interesting bioactivities including anti-inflamma-

tory (11, 13, 16), antiplatelet-aggregatory (11, 16–18), cytotoxic (10, 11, 16, 19, 20), and antitumor activity (9, 22) as well as activity against bacteria and mycoplasma (21). Falcarinol appears to be the most bioactive among polyacetylenes of the falcarinol-type. The bioactivity of falcarinol is probably associated with its hydrophobicity and its ability to form an extremely stable carbocation with the loss of water, thereby acting as a very reactive alkylating agent toward various biomolecules (9, 11). This is supported by the fact that falcarinol is a strong contact allergen (23). A similar mode of action is expected for other falcarinol-type polyacetylenes with a hydroxyl group at C-3 (9, 11).

The beneficial effects of polyacetylenes of the falcarinol-type occur at nontoxic concentrations and thus represent pharmacologically useful properties indicating that polyacetylenes may be important nutraceuticals of vegetables. In the human diet, carrots are the major dietary source of falcarinol-type polyacetylenes, in particular falcarinol and falcarindiol (8, 13, 14). It has been demonstrated that falcarinol but not β -carotene could stimulate differentiation of primary mammalian cells of mammary origin in concentrations between 1 ng/mL and 10 ng/mL falcarinol. Toxic effects were found > 50 ng/mL falcarinol, while β -carotene had

*To whom correspondence should be addressed. Tel: +45 89 99 15 56. Fax: +45 89 99 11 66. E-mail: stig.purup@agrsci.dk.




Figure 1. Chemical structures of the polyacetylenes falcarinol, falcarindiol, panaxydol, and falcarinon tested on human intestinal epithelial cells of normal (FHs 74 Int.) and cancer (Caco-2) origin in the present investigation.

no effect even at 100 $\mu\text{g/mL}$ (8). This biphasic effect (hormesis) on cell proliferation has also recently been demonstrated for falcarindiol (24). This is fully in accordance with the hypothesis that toxic compounds have beneficial effects at certain lower concentrations (25). Therefore, falcarinol-type polyacetylenes appear to be the bioactive components in carrots that could explain their health promoting properties, rather than carotenoids, polyphenols, or other types of primary and/or secondary metabolites. This is also in accordance with a recent study by Metzger et al. (13) who found that the anti-inflammatory activities of purple carrot cultivars was not due to the high levels of polyphenols but instead to their content of falcarinol-type polyacetylenes. Studies on the bioavailability of particular falcarinol in humans support the hypothesis that aliphatic C_{17} -polyacetylenes of the falcarinol-type may play an important role in human health (7, 11). When falcarinol was administered orally via carrot juice (13.3 mg falcarinol/L carrot juice) in amounts of 900 mL, it was rapidly absorbed, reaching a maximum concentration in serum of 2.5 ng/mL at 2 h after dosing (7, 11). This is within the range where the in vitro data indicate a potentially beneficial physiological effect and a possible inhibitory effect on the proliferation of cancer cells (8, 12).

The potential anticancer activity of carrots and falcarinol has been demonstrated in an established rat model for colon cancer by injections of the carcinogen azoxymethane in the inbred rat strain BDXI (9). The carrot and falcarinol treatments showed a significant tendency to reduce the number of (pre)cancerous lesions (aberrant crypt foci) with increasing size of lesions in physiologically relevant concentrations, i.e., in amounts corresponding to a daily human consumption of 400–600 g fresh weight of carrot (9). In this study, the effect of the falcarinol diet was shown to have a larger effect on pre(cancerous) lesions compared to that in the carrot diet, although the differences between the falcarinol and the carrot diet was not significant. This could indicate that other compounds in carrots may interact with falcarinol thereby affecting its effectiveness in vivo, in particular its anticancer effect. This raises some interesting questions regarding the interaction between polyacetylenes and other metabolites in carrots. For example, is the anticancer effect of falcarinol affected by other carrot polyacetylenes such as falcarindiol, and if yes, how does this interaction most likely affect the bioactivity of falcarinol? Alternatively, is the bioactivity of carrot polyacetylenes in vivo affected by the presence of other metabolites in carrots?

In an attempt to answer these questions, the differential effect of common polyacetylenes of the falcarinol-type (falcarinol, falcarindiol, and panaxydol) as well as different combinations of falcarinol and falcarindiol and carrot extracts with different polyacetylene profiles on the proliferation of human intestinal epithelial cells of normal (FHs 74 Int.) and cancer (Caco-2) origin were investigated. The hypothesis that a hydroxyl group at C-3 is important for the bioactivity of falcarinol-type polyacetylenes was investigated by comparing the growth inhibitory effect of falcarinol and the oxidized form of falcarinol, falcarinon, on intestinal cell proliferation.

MATERIALS AND METHODS

Chemicals and Materials. Ethyl acetate (EtOAc), ethanol (EtOH, 96%), and *n*-hexane of high-performance liquid chromatography (HPLC) grade (99.9% HPLC grade) and diethyl ether (Et_2O ; 99.9% p.a. for GC) were obtained from Sigma-Aldrich (Steinheim, Germany). Activated manganese(IV) oxide (MnO_2 ; ReagentPlus $\geq 99\%$) and anhydrous sodium sulfate (Na_2SO_4) were obtained from Sigma-Aldrich. Preparative TLC plates (20 \times 20 cm, silica gel 60 F₂₅₄, 2.5 mm thickness) were obtained from Merck (Darmstadt, Germany).

Polyacetylene Standards. Authentic standards of the polyacetylenes (3*R*)-falcarinol and (3*R*, 8*S*)-falcarindiol (Figure 1) were isolated from 2.5 kg of carrot roots (*Daucus carota*, cv. Bolero) and (3*R*)-panaxydol and (3*R*)-falcarinol (Figure 1) from 2 kg of American ginseng (*Panax quinquefolium*) roots according to the procedure described by Kidmose et al. (14) and Christensen et al. (15), respectively. Falcarinol, falcarindiol, and panaxydol were obtained in a purity $> 98\%$, which was demonstrated by analyzing the isolated polyacetylenes by reverse phase (RP) HPLC–photodiode array detection (PAD) according to the method described by Christensen and Kreutzmann (34). Polyacetylenes were identified by UV, mass spectrometry (MS) [gas chromatography (GC)–MS (EI, 70 eV)], NMR (^1H and ^{13}C NMR, and ^1H – ^1H and ^1H – ^{13}C correlation spectroscopy recorded in CDCl_3 with TMS as internal standard), and optical rotation, and the complete spectral data set corresponded fully with literature values for falcarinol (9, 22, 26–28), falcarindiol (22, 27–29), and panaxydol (19, 30–32), respectively.

Falcarinon (Figure 1) was produced by the oxidation of falcarinol by activated MnO_2 (33). Activated MnO_2 (100 mg) was added to a solution of falcarinol (25 mg) in Et_2O (50 mL) and the mixture stirred at room temperature for 2 h under dark conditions. The reaction mixture was filtered and the residue washed with Et_2O . The filtrate was concentrated in vacuo (30 $^\circ\text{C}$) in dim light and the residue purified by preparative TLC (*n*-hexane– Et_2O , 3:1) to give 18 mg of falcarinon as a pale yellow oil in a purity $> 98\%$ as shown by RP-HPLC–PAD (34). UV λ_{max} (Et_2O): 228, 262, 276, 292 nm. GC–MS (EI, 70 eV): m/z 242 [$\text{M}]^+$ (9), 185 (5), 171 (11), 157 (57), 145 (36), 143 (20), 131 (63), 129 (80), 128 (45), 117 (100), 115 (83), 103 (87), 91 (59), 77 (76), 55 (94). ^1H NMR (300 MHz, CDCl_3 , TMS as internal standard): δ 0.88 (3H, t, 7.1 Hz, H-17), 1.28 (8H, m, H-13–H-16), 1.37 (2H, m, H-12), 2.04 (2H, dt, 7.0, 7.5 Hz, H-11), 3.12 (2H, d, 6.9 Hz, H-8), 5.38 (1H, m, H-9), 5.57 (1H, m, dtt, 1.8, 7.5, 10.5 Hz, H-10), 6.19 (1H, dd, 0.9, 10.1 Hz, H-1), 6.40 (1H, dd, 10.1, 17.4 Hz, H-1), 6.55 (1H, dd, 0.9, 17.4 Hz, H-2). ^{13}C NMR (75 MHz, CDCl_3 , TMS as internal standard): δ 14.1 (CH_3 , C-17), 18.0 (CH_2 , C-8), 22.6 (CH_2 , C-16), 27.3 (CH_2 , C-11), 29.1 (CH_2 , C-14), 29.2 (CH_2 , C-13), 29.2 (CH_2 , C-12), 31.8 (CH_2 , C-15), 63.6 (C, C-6), 70.6 (C, C-5), 77.1 (C, C-7), 88.0 (C, C-4), 120.7 (CH, C-9), 133.9 (CH, C-10), 134.1 (CH_2 , C-1), 137.8 (CH, C-2), 177.8 (C, C-3).

Carrot Root Extracts for Testing. Frozen carrot shreds (20 g) from the cultivars cv. Bolero (carrot extract 1) and cv. Purple Haze (carrot extract 2) were homogenized to a puree using an Ultra-Turrax T25 for 1 min followed by extraction with 60 mL of EtOAc overnight under stirring at room temperature. Before stirring, the samples were exposed to ultrasound for 10 min. After the first extraction, the samples were extracted with an additional 60 mL of EtOAc for 3 h under stirring at room temperature. The EtOAc extracts were combined and the water removed from the combined extract by adding anhydrous Na_2SO_4 (15 g) followed by filtration through a Munktell filter paper, grade 3W. The dried extract was evaporated in vacuo at 25 $^\circ\text{C}$ in dim light, and the residue obtained was dissolved in EtOH (5 mL) and filtered through a 0.45 μm




Figure 2. Effect of increasing concentrations of faltarinol, faltarindiol, and panaxydol on cell proliferation in (A,B) Caco-2 and (C,D) FHs 74 Int. cells cultured for 72 h in either (A,C) 0.625% FCS or (B,D) 10% FCS. Values are least-squares means obtained from cultures in quadruple samples from two experiments and presented as relative to proliferation obtained in medium without polyacetylenes (basal medium). Standard error of the mean (SEM) values obtained in the ANOVA analysis were 0.15, 0.08, 0.06, and 0.04 (relative values) for A, B, C, and D, respectively. Values significantly different from proliferation obtained in basal medium are indicated: c, $P < 0.05$; b, $P < 0.01$; a, $P < 0.001$.

Nylon, Cameo filter (MSI, Westborough, MA) and analyzed by RP-HPLC-PAD for faltarinol, faltarindiol, and faltarindiol 3-acetate according to the method described by Christensen and Kreutzmann (34).

Cell Culture. Caco-2 cells (a kind gift from Dr. Jan Trige Rasmussen, Protein Chemistry Laboratory, University of Aarhus, Denmark) were cultured in 96-well plates at a density of 4000 cells/well in Dulbecco's minimal essential medium (DMEM) (Gibco 11960-044), supplemented with fetal calf serum (FCS, 10%; Cambrex Bio Science, Copenhagen, Denmark), *N*-(2-hydroxyethyl)piperazine-*N'*-ethanesulfonic acid (HEPES) (10 mM; Invitrogen A/S, Tåstrup, Denmark), Glutamax (2 mM; Invitrogen A/S), and antibiotic solution (P-3539; Sigma-Aldrich A/S, Vællensbæk Strand, Denmark) containing penicillin and streptomycin for 24 h (pretreatment period). FHs 74 Int. cells (ATCC; purchased from LGC Nordic AB, Borås, Sweden) were cultured in 96-well plates at a density of 1500 cells/well in DMEM (Gibco 42430-025), supplemented with fetal calf serum (FCS, 10%; Cambrex Bio Science, Copenhagen, Denmark), HEPES (10 mM; Invitrogen A/S, Tåstrup, Denmark), bovine insulin (7.5 mg/L; Sigma-Aldrich A/S), human recombinant EGF (30 μ g/L; Austral Biologicals, San Ramon CA, USA), NCTC-135 with glutamin (8%; Invitrogen A/S), nonessential amino acids (1%; Invitrogen A/S), sodium puruvate (0.4 mM; Invitrogen A/S), oxaloacetate (0.1 g/L; Sigma-Aldrich A/S), Glutamax (2 mM; Invitrogen A/S), and antibiotic solution (P-0781; Sigma-Aldrich A/S) containing penicillin and streptomycin for 24 h (pretreatment period). At 60 to 80% confluence, cells were washed carefully with phosphate buffered saline (PBS) (Invitrogen A/S), and treatment medium (200 μ L per well in 96-well plates) was added to individual wells for a total of 72 h, with medium changed after 48 h. For each cell line, growth curves were obtained with different numbers of cells and different concentrations of FCS for 8–10 successive days of culture. In this way, we ensured that cells were in the sigmoid phase of the growth curve when components to be tested were added to the culture medium.

Treatment medium contained faltarinol, faltarindiol, or panaxydol in different concentrations (0.001, 0.01, 0.1, 1, 2.5, 5, 10, and 20 μ g/mL), faltarinol or faltarinol in different concentrations (1, 2.5, 5, 10, and 20 μ g/mL), or carrot extracts (0.01, 0.05, 0.1, 0.5, and 1% in culture medium). These treatment media were prepared from stock concentrations of polyacetylenes in 96% EtOH or carrot extracts resuspended in 96% EtOH. Concentrations of EtOH in culture media never exceeded 1%. Experiments also included studies of potential synergistic and/or antagonistic effects of quantitative major and most bioactive polyacetylenes in carrots. For these studies, faltarinol and faltarindiol were added to culture medium in combinations. Keeping one of the polyacetylenes constant at 10 ng/mL or 1 μ g/mL, the two compounds were tested for effects in the ratios 1:1, 1:5, 1:10, 5:1, and 10:1. Treatment medium furthermore contained FCS (Cambrex Bio Science, Copenhagen, Denmark) in concentrations of either 0.625% or 10%. Cell proliferation was determined by a resazurin metabolism assay (AlamarBlue, BioSource, AH diagnostics, Aarhus, Denmark). Cell culture medium was carefully aspirated and the AlamarBlue reagent was added to all wells and measured after 2 h incubation according to the manufacturer.

Statistics. Statistical analyses were performed using the general linear models procedure (PROC GLM) of SAS Institute Inc. Different concentrations of the three polyacetylenes (Figure 2) were tested using a model including the systematic effects of the source of polyacetylenes ($n = 3$), concentrations of polyacetylenes (nine levels), and their interaction. The effect of the assay ($n = 2$) was included as a blocking factor. Different concentrations of faltarinol and faltarinol (Figure 3) were tested using a model including the systematic effect of polyacetylenes ($n = 2$), concentration of polyacetylenes (five levels), and their interaction. The effect of the assay ($n = 2$) was included as a blocking factor. Different concentrations of the two carrot extracts (Figure 4) were tested using a model




Figure 3. Effect of increasing concentrations of falcariinol and falcarinon on cell proliferation in (A) Caco-2 and (B) FHs 74 Int. cells cultured for 72 h in 0.625% FCS. Values are least-squares means obtained from cultures in quadruple samples from two experiments and presented as relative to proliferation obtained in medium without falcariinol (basal medium). SEM values obtained from ANOVA analysis were 6.4 and 15.3 (relative values) for A and B, respectively. Values significantly different from proliferation obtained in basal medium are indicated: b, $P < 0.01$; a, $P < 0.001$. Values showing a tendency to be different from proliferation obtained in basal medium are indicated: †, $P < 0.1$.

including the systematic effect of extract ($n = 2$), concentration of extract (five levels), and their interaction. The effect of the assay ($n = 2$) was included as a blocking factor. The residual mean error was used as the error term for F-tests. All results shown in Figures 2–4 are least-squares means obtained from cultures in quadruple samples from two experiments and presented as relative to proliferation obtained in basal medium.

RESULTS AND DISCUSSION

The effects of falcariinol-type polyacetylenes and carrot extracts were evaluated in two different intestinal cell lines of normal and cancer origin, respectively. We have previously used both of these intestinal epithelial cell lines for identification of bioactive components in complex natural mixtures such as milk (35, 36). To be able to compare results obtained from two different cell lines, growth curves were performed for each cell line to ensure that cells were in the sigmoid phase of the growth curve when components to be investigated were added to the culture medium. Therefore, the initial number of cells was different for the two individual cell lines.

Effect of Polyacetylenes on Intestinal Cell Proliferation. The effects of the polyacetylenes, falcariinol, falcariindiol, and panaxydol were first evaluated in the human intestinal epithelial cell line of cancer origin, Caco-2, using a resazurin metabolism assay as an index of cell proliferation. Treatment of the cells with concentrations ranging from 1 ng/mL to 20 µg/mL showed a hormesis effect of all three polyacetylenes. Falcariinol, falcariindiol, and panaxydol in a concentration of 100 ng/mL increased cell proliferation




Figure 4. Effect of increasing concentrations of carrot extracts on cell proliferation in (A) Caco-2 and (B) FHs 74 Int. cells cultured for 72 h in 0.625% FCS. Concentrations correspond to percentages of the stock solutions given in Table 2. Values are least-squares means obtained from cultures in quadruple samples from two experiments and presented as relative to proliferation obtained in medium without extracts (basal medium). SEM values obtained in the ANOVA analysis were 2.8 and 7.5 (relative values) for A and B, respectively. Values significantly different from proliferation obtained in basal medium are indicated: c, $P < 0.05$; a, $P < 0.001$. Carrot extract 1 contained approximately 50% of the amount of falcariinol present in carrot extract 2, but approximately 4 and 2 times more of falcariindiol and falcariindiol 3-acetate than in carrot extract 2.

by 30, 42, and 60% under basal growth conditions (0.625% FCS) ($P < 0.05$; Figure 2A). However, in the presence of 10% FCS, no significant increase in cell proliferation was observed with the three polyacetylenes (Figure 2B). Treatment with falcariinol at a concentration of 2.5 µg/mL, and falcariindiol and panaxydol at concentrations of 10 µg/mL significantly reduced proliferation under basal growth conditions ($P < 0.05$; Figure 2A). In the presence of 10% FCS, concentrations of 5, 20, and 10 µg/mL of falcariinol, falcariindiol, and panaxydol, respectively, were needed to reduce proliferation significantly ($P < 0.01$; Figure 2B). The relative inhibitory potency was falcariinol > panaxydol > falcariindiol with concentrations corresponding to 50% inhibition of basal response of 2.5–5 µg/mL, 5–10 µg/mL, and 10–20 µg/mL, respectively. Maximal inhibition at 20 µg/mL corresponded to approximately 80% inhibition of cell proliferation under basal growth conditions (Figure 2A). Under FCS stimulated growth conditions, concentrations corresponding to 50% inhibition were 5–10 µg/mL, 10–20 µg/mL and > 20 µg/mL, for falcariinol, panaxydol, and falcariindiol, respectively. Maximal inhibition at 20 µg/mL corresponded to approximately 90, 65, and 30% for falcariinol, panaxydol, and falcariindiol, respectively (Figure 2B).

We also determined whether falcariinol, falcariindiol, and panaxydol had similar effects in the normal human intestinal epithelial cell line, FHs 74 Int. In these cells, no hormesis effect was observed. Instead, a significantly inhibitory effect on proliferation was seen with concentrations of falcariinol, falcariindiol, and panaxydol at 5, 10, and 2.5 µg/mL, respectively ($P < 0.001$; Figure 2C and D) in the presence of both 0.625 and 10% FCS,

except for faltarindiol which had no inhibitory effect in the presence of 10% FCS. The relative inhibitory potency was panaxydol > faltarinol > faltarindiol with concentrations corresponding to 50% inhibition of basal response of 1–2.5 $\mu\text{g/mL}$, 2.5–5 $\mu\text{g/mL}$, and 10–20 $\mu\text{g/mL}$, respectively. When 10% FCS was present in the culture medium, concentrations corresponding to 50% inhibition were 1–2.5 $\mu\text{g/mL}$ and 5–10 $\mu\text{g/mL}$ for panaxydol and faltarinol, respectively (**Figure 2C and D**).

The present study shows that faltarinol, faltarindiol, and panaxydol affect cell proliferation of human intestinal cells in a dose-dependent way and that the presence of FCS in culture medium also influences these responses. In a recent study, Caco-2 cells were chosen as the *in vitro* model for assessment of the bioactivity of faltarinol, but faltarindiol and panaxydol were not examined (12). However, the effect of faltarinol was only studied in the presence of 10% FCS in the culture medium. Such high concentration of FCS in culture medium may protect intestinal cells from growth-inhibitory compounds and furthermore may include relatively high concentrations of compounds that affect the bioactivity of the tested compounds. This may explain why the hormesis effect of polyacetylenes obtained in Caco-2 cells at 0.625% FCS could not be seen in the presence of 10% FCS and that higher concentrations of polyacetylenes were needed to inhibit cell proliferation (**Figure 2**). These results indicate that FCS in high concentrations in culture medium protects intestinal cells against the inhibitory effects of the tested polyacetylenes. On the basis of these observations, we recommend testing the bioactivity of faltarinol-type polyacetylenes in Caco-2 cells and similar cell models in low concentrations of FCS.

The inhibitory potency was clearly different between the tested polyacetylenes in the cancer and normal intestinal cells. In Caco-2 cells, faltarinol was the most potent inhibitor of cell proliferation, while in FHs cells, panaxydol was more potent than faltarinol. The effect of faltarindiol was similar in both cell lines, although its effect was much lower with 50% inhibition of basal response at 10–20 $\mu\text{g/mL}$ (**Figure 2**). The present results confirm the results of other studies that faltarinol and panaxydol are much more cytotoxic than faltarindiol and similar polyacetylenes of the faltarinol-type such as faltarindiol 3-acetate and 8-*O*-methylfaltarindiol (10). Faltarindiol has, for example, the ability to generate two active centers for nucleophilic attack, which reduces the lipophilic character of this compound and hence its reactivity, which could explain the lower reactivity of this compound compared to that of faltarinol and panaxydol. The lower reactivity and potential anticancer effect of faltarindiol compared to that of faltarinol are also in accordance with the observed nonallergenic properties of faltarindiol (23). The inhibitory activity of the tested polyacetylenes toward Caco-2 cells therefore confirm that the anticancer activity of polyacetylenes of the faltarinol-type is associated with their ability to form extremely stable carbocations, thereby acting as very reactive alkylating agents toward biomolecules.

Effect of Oxidation of Faltarinol on Intestinal Cell Proliferation.

The importance of a hydroxyl group in the C-3 position of faltarinol was investigated by testing the inhibitory effect of the oxidized form of faltarinol, faltarinon, on intestinal cell proliferation. Faltarinol and faltarinon, were added to Caco-2 and FHs 74 Int. cells for comparison of the effects on cell proliferation. In accordance with the results reported above, faltarinol significantly inhibited cell proliferation in Caco-2 cells at 2.5 $\mu\text{g/mL}$ ($P < 0.01$) and in FHs 74 Int. cells at 5 $\mu\text{g/mL}$ ($P < 0.01$) (**Figure 3**). Faltarinon, however, only inhibited proliferation in Caco-2 cells at the concentration of 20 $\mu\text{g/mL}$ ($P < 0.001$) and only tended to decrease proliferation in FHs 74 Int. cells at this concentration ($P < 0.10$). Furthermore, the dose of 20 $\mu\text{g/mL}$

Table 1. Effects of Different Ratios of Faltarinol and Faltarindiol on Caco-2 and FHs 74 Int. Cells in Culture Medium Containing 0.625% FCS^a

		faltarinol ($\mu\text{g/mL}$)			
		0	1	5	10
Caco-2 cells					
faltarindiol ($\mu\text{g/mL}$)	0	1.0	1.01 \pm 0.25	0.36 \pm 0.25	0.18 \pm 0.01
	1	1.35 \pm 0.36	0.71 \pm 0.08 ^b	0.58 \pm 0.04	0.22 \pm 0.08
	5	1.03 \pm 0.32	0.53 \pm 0.06 ^b	nd ^c	nd
	10	0.52 \pm 0.17	0.17 \pm 0.03 ^b	nd	nd
FHs 74 Int. cells					
faltarindiol ($\mu\text{g/mL}$)	0	1.0	1.02 \pm 0.06	0.32 \pm 0.35	0.06 \pm 0.01
	1	0.92 \pm 0.01	0.51 \pm 0.30 ^b	0.47 \pm 0.01	0.36 \pm 0.28
	5	0.90 \pm 0.04	0.34 \pm 0.24 ^b	nd	nd
	10	0.60 \pm 0.14	0.14 \pm 0.13 ^b	nd	nd

^a Cells were cultured for 72 h, and cell proliferation was determined by a resazurin metabolism assay. Mean \pm standard deviation is shown for each combination of the two tested polyacetylenes. Data are presented relative to cell proliferation obtained in medium without polyacetylenes for two individual experiments. ^b Significant synergistic inhibitory effect on cell proliferation compared to single-compound assay ($P < 0.01$). ^c nd: not determined.

only caused 40–45% reduction in cell proliferation, and there were no differences between normal and cancer cells (**Figure 3**). These results demonstrate that the oxidized form of faltarinol, faltarinon, is a much less potent inhibitor of cell proliferation in intestinal cells of both normal and cancer origin and strongly support the hypothesis that the mode of action of faltarinol-type polyacetylenes is due to their ability to lose water through the hydroxyl group at C-3, thereby generating reactive carbocations. This mode of action of faltarinol-type polyacetylenes makes it reasonable to assume that these compounds can affect each other's anticancer activity in an antagonistic and/or synergistic manner. Consequently, the inhibitory effect of faltarinol and faltarindiol on Caco-2 and FHs 74 Int. cells was tested in different concentration ratios. Keeping one of the polyacetylenes constant at either 10 ng/mL or 1 $\mu\text{g/mL}$, Caco-2 and FHs 74 Int. cells were incubated with faltarinol and faltarindiol in the ratios 1:1, 1:5, 1:10, 5:1, and 10:1. These studies were again performed in a medium containing 0.625% FCS or 10% FCS. When a concentration of 10 ng/mL of faltarinol or faltarindiol was added to Caco-2 or FHs 74 Int. cells at either 0.625% or 10% FCS present in the medium, increasing the concentration of one or the other polyacetylene showed no difference from what would have been expected from the single-compound assay (data not shown). However, at a concentration of 1 $\mu\text{g/mL}$ of faltarinol, a synergistic response for the inhibitory effect of cell proliferation was observed by adding faltarindiol in 1, 5, and 10 times the concentration of faltarinol (**Table 1**) for both Caco-2 and FHs 74 Int. cells but only with 0.625% FCS present in the medium. By keeping faltarindiol constant, no synergistic response of cell proliferation was observed by adding faltarinol in 5 and 10 times the concentration of faltarindiol in Caco-2 nor FHs 74 Int. cells (**Table 1**).

The results of these studies demonstrate that faltarindiol in low doses can have very potent inhibitory effects on intestinal cell proliferation when found in combination with low doses of faltarinol (**Table 1**). Synergistic interactions could therefore be an important factor in relation to the anticancer activity of faltarinol-type polyacetylenes, although this activity clearly depends on the concentration and the ratio of the compounds.

Effect of Carrot Extracts on Intestinal Cell Proliferation. To further determine the effect of polyacetylenes in combinations and hence their possible synergistic interactions, the effects of two

Table 2. Concentrations of Falcarinol, Falcarindiol, and Falcarindiol 3-Acetate in EtOAc Extracts from the Carrot Cultivars cv. Bolero (Carrot Extract 1) and cv. Purple Haze (Carrot Extract 2)^a

carrot extract	falcarinol ($\mu\text{g/mL}$)	falcarindiol ($\mu\text{g/mL}$)	falcarindiol 3-acetate ($\mu\text{g/mL}$)
1	281.1	596.0	195.8
2	645.7	152.3	107.1

^a Concentrations are given in stock solutions of which different concentrations were prepared in cell culture medium.

carrot EtOAc extracts were studied in Caco-2 and FHs 74 Int. cells. Concentrations of falcarinol, falcarindiol, and falcarindiol 3-acetate in carrot extracts are shown in **Table 2**. Carrot extract 1 contained approximately 50% of the falcarinol in carrot extract 2, but approximately 4 and 2 times more falcarindiol and falcarindiol 3-acetate, respectively, than carrot extract 2. Both extracts had significant inhibitory effects on intestinal cell proliferation in the highest concentration of 0.5 and 1% of extract in culture medium (**Figure 4**). In Caco-2 cells, carrot extract 2 tended ($P < 0.10$) to be a more potent inhibitor of proliferation than carrot extract 1, as significant inhibition was obtained also with 0.1% extract giving an approximate 20% decrease in proliferation (**Figure 4A**). In FHs 74 Int. cells, there was no difference between the effects of the two extracts (**Figure 4B**), but the effect at a given concentration was larger than that in Caco-2 cells. At the highest concentrations of extracts, cell proliferation was reduced to 6% of proliferation in FHs 74 Int. cells, while in Caco-2 cells, cell proliferation was reduced to only 36% (**Figure 4**).

The more potent inhibitory activity of carrot extract 2 than of carrot extract 1 in Caco-2 cells (**Figure 4A**) is in accordance with the approximately 3 times higher concentration of falcarinol in carrot extract 2, as falcarinol is a more potent inhibitor than falcarindiol (**Figure 2**). Comparing the effects of the carrot extracts with the effects of falcarinol or falcarindiol alone, we showed that the mixture of polyacetylenes found in the extracts was more potent in inhibiting cell proliferation. As an example, the 1% carrot extract 1 prepared from a stock solution by diluting 100 times in cell culture medium (**Table 2**) contained falcarinol, falcarindiol, and falcarindiol 3-acetate in concentrations of 2.81, 5.96, and 1.96 $\mu\text{g/mL}$, respectively, and caused a reduction in cell proliferation to 36%. For comparison, a concentration of 2.5 $\mu\text{g/mL}$ of falcarinol decreased proliferation to 68%, while falcarindiol caused no decrease in cell proliferation (**Figures 2 and 4**, and **Table 2**). Concentrations in falcarinol of 5 $\mu\text{g/mL}$ or falcarindiol of 10–20 $\mu\text{g/mL}$ were required to obtain a decrease in cell proliferation corresponding to 36%. However, addition of carrot extract 2 to Caco-2 cells did not decrease cell proliferation more than carrot extract 1 despite a concentration in falcarinol of 6.46 $\mu\text{g/mL}$. These studies suggest that falcarinol in combination with the relatively high concentrations of falcarindiol and falcarindiol 3-acetate in carrot extract 1 has a synergistic effect on Caco-2 cell proliferation. In FHs 74 Int. cells, a similar synergistic effect can be suggested as carrot extract 1 or 2 both caused a decrease in cell proliferation to 30 and 6% in concentrations of 0.5% and 1%, respectively, of the extracts. To obtain such decreases in cell proliferation of FHs 74 Int. cells by falcarinol alone, a concentration of 10 $\mu\text{g/mL}$ would be necessary.

The synergistic effect observed between falcarinol and other falcarinol-type polyacetylenes in carrots are contradictory to the anticancer effect previously observed for a carrot diet compared to a falcarinol diet in rats with induced (pre)cancerous lesions (9). The present results therefore clearly suggest that other metabolites in carrots to some extent may interact with falcarinol and other polyacetylenes in carrots, thereby affecting their effectiveness in vivo.

ACKNOWLEDGMENT

We thank A. K. Nielsen and K. B. Poulsen for their skilled technical assistance.

Note Added After ASAP Publication

The original ASAP publication of August 20, 2009, did not reflect all galley proof corrections. These have been incorporated in the publication of September 1, 2009.

LITERATURE CITED

- (1) Steinmetz, K. A.; Potter, J. D. Vegetables, fruit, and cancer prevention: A review. *J. Am. Diet. Assoc.* **1996**, *96*, 1027–1039.
- (2) Block, G.; Patterson, B.; Subar, A. Fruit, vegetables, and cancer prevention: A review of the epidemiological evidence. *Nutr. Cancer* **1992**, *18*, 1–29.
- (3) Greenvald, P.; Clifford, C. K.; Milner, J. A. Diet and cancer prevention. *Eur. J. Cancer* **2001**, *37*, 948–965.
- (4) Kris-Etherton, P. M.; Etherton, T. D.; Carlson, J.; Gardner, C. Recent discoveries in inclusive food-based approaches and dietary patterns for reduction in risk for cardiovascular disease. *Curr. Opin. Lipid* **2002**, *13*, 397–407.
- (5) Maynard, M.; Gunnell, D.; Emmett, P.; Frankel, S.; Smith, G. D. Fruit, vegetables, and antioxidants in childhood and risk of adult cancer: the Boyd Orr cohort. *J. Epidemiol. Community Health* **2003**, *57*, 218–225.
- (6) Trichopoulou, A.; Naska, A.; Antoniou, A.; Friel, S.; Trygg, K.; Turrini, A. Vegetable and fruit: the evidence in their favour and the public health perspective. *Int. J. Vitam. Nutr. Res.* **2003**, *73*, 63–69.
- (7) Brandt, K.; Christensen, L. P.; Hansen-Møller, J.; Hansen, S. L.; Haraldsdóttir, J.; Jespersen, L.; Purup, S.; Kharazmi, A.; Barkholdt, V.; Frøkiær, H.; Kobæk-Larsen, M. Health promoting compounds in vegetables and fruits: A systematic approach for identifying plant components with impact on human health. *Trends Food Sci. Technol.* **2004**, *15*, 384–393.
- (8) Hansen, S. L.; Purup, S.; Christensen, L. P. Bioactivity of falcarinol and the influence of processing and storage on its content in carrots (*Daucus carota* L.). *J. Sci. Food Agric.* **2003**, *83*, 1010–1017.
- (9) Kobæk-Larsen, M.; Christensen, L. P.; Vach, W.; Ritskes-Hoitinga, J.; Brandt, K. Inhibitory effects of feeding with carrots or (–)-falcarinol on development of azoxymethane-Induced preneoplastic lesions in the rat colon. *J. Agric. Food Chem.* **2005**, *53*, 1823–1827.
- (10) Zidorn, C.; Johrer, K.; Ganzera, M.; Schubert, B.; Sigmund, E. M.; Mader, J.; Greil, R.; Ellmerer, E. P.; Stuppner, H. Polyacetylenes from the Apiaceae vegetables carrot, celery, fennel, parsley, and parsnip and their cytotoxic activities. *J. Agric. Food Chem.* **2005**, *53*, 2518–2523.
- (11) Christensen, L. P.; Brandt, K. Bioactive polyacetylenes in food plants of the Apiaceae family: occurrence, bioactivity and analysis. *J. Pharm. Biomed. Anal.* **2006**, *41*, 683–693.
- (12) Young, J. F.; Duthie, S. J.; Milne, L.; Christensen, L. P.; Duthie, G. G.; Bestwick, C. S. Biphasic effect of falcarinol on CaCo-2 cell proliferation, DNA damage, and apoptosis. *J. Agric. Food Chem.* **2007**, *55*, 618–623.
- (13) Metzger, B. T.; Barnes, D. M.; Reed, J. D. Purple carrot (*Daucus carota* L.) polyacetylenes decrease lipopolysaccharide-induced expression of inflammatory proteins in macrophage and endothelial cells. *J. Agric. Food Chem.* **2008**, *56*, 3554–3560.
- (14) Kidmose, U.; Hansen, S. L.; Christensen, L. P.; Edelenbos, M.; Nørbæk, R. Effects of genotype, processing, storage, and root size on bioactive compounds (carotenoids, polyacetylenes, 6-methoxymellein) in organically grown carrots (*Daucus carota* L.). *J. Food Sci.* **2004**, *69*, S388–S394.
- (15) Christensen, L. P.; Jensen, M.; Kidmose, U. Simultaneous determination of ginsenosides and polyacetylenes in American ginseng root (*Panax quinquefolium* L.) by high-performance liquid chromatography. *J. Agric. Food Chem.* **2006**, *54*, 8995–9003.
- (16) Christensen, L. P.; Jakobsen, H. B. Polyacetylenes: Distribution in Higher Plants, Pharmacological Effects and Analysis. In *Thin Layer Chromatography in Phytochemistry*; Waksmundzka-Hajnos, M.,

- Sherma, J., Kowalska, T., Eds.; Chromatographic Science Series; CRC Press/Taylor & Francis Group: Boca Raton, FL, 2008; Vol. 99, pp 757–816.
- (17) Appendino, G.; Tagliapietra, S.; Nano, G. M.; Picci, V. An anti-platelet acetylene from the leaves of *Ferula communis*. *Fitoterapia* **1993**, *64*, 179.
 - (18) Alanko, J.; Kurahashi, Y.; Yoshimoto, T.; Yamamoto, S.; Baba, K. Panaxynol, a polyacetylene compound isolated from oriental medicines, inhibits mammalian lipoxygenases. *Biochem. Pharmacol.* **1994**, *48*, 1979–1981.
 - (19) Fujimoto, Y.; Satoh, M.; Takeuchi, N.; Kirisawa, M. Cytotoxic acetylenes from *Panax quinquefolium*. *Chem. Pharm. Bull.* **1991**, *39*, 521–523.
 - (20) Matsunaga, H.; Katano, M.; Yamamoto, H.; Fujito, H.; Mori, M.; Takata, K. Cytotoxic activity of polyacetylene compounds in *Panax ginseng* C. A. Meyer. *Chem. Pharm. Bull.* **1990**, *38*, 3480–3482.
 - (21) Kobaisy, M.; Abramowski, Z.; Lermer, L.; Saxena, G.; Hancock, R. E. W.; Towers, G. H. N. Antimycobacterial polyynes of Devil's Club (*Oplopanax horridus*), a North American native medicinal plant. *J. Nat. Prod.* **1997**, *60*, 1210–1213.
 - (22) Bernart, M. W.; Cardellina, J. H. II; Balaschak, M. S.; Alexander, M.; Shiemaker, R. H.; Boyd, M. R. Cytotoxic falcarinol oxylipins from *Dendropanax arboreus*. *J. Nat. Prod.* **1996**, *59*, 748–753.
 - (23) Hansen, L.; Hammershøj, O.; Boll, P. M. Allergic contact dermatitis from falcarinol isolated from *Schefflera arboricola*. *Contact Dermatitis* **1986**, *14*, 91–93.
 - (24) Young, J. F.; Christensen, L. P.; Theil, P. K.; Oksbjerg, N. The polyacetylenes falcarinol and falcarindiol affect stress responses in myotube cultures in a biphasic manner. *Dose–Response* **2008**, *6*, 239–251.
 - (25) Calabrese, E. J.; Baldwin, L. A. The hormetic dose-response model is more common than the threshold model in toxicology. *Toxicol. Sci.* **2003**, *71*, 246–250.
 - (26) Hirakura, K.; Morita, M.; Nakajima, K.; Ikeya, Y.; Mitsunashi, H. Three acetylenic compounds from roots of *Panax ginseng*. *Phytochemistry* **1992**, *31*, 899–903.
 - (27) Czepa, A.; Hofmann, T. Structural and sensory characterization of compounds contributing to the bitter off-taste of carrots (*Daucus carota* L.) and carrot puree. *J. Agric. Food Chem.* **2003**, *51*, 3865–873.
 - (28) Nitz, S.; Spraul, M. H.; Drawert, F. C₁₇ Polyacetylenic alcohols as the major constituents in roots of *Petroselinum crispum* Mill. ssp. *tuberosum*. *J. Agric. Food Chem.* **1990**, *38*, 1445–1447.
 - (29) Lemmich, E. Constituents of Umbelliferous plants. 23. The absolute-configuration of the acetylenic-compound falcarindiol. *Phytochemistry* **1981**, *20*, 1419–1420.
 - (30) Hirakura, K.; Morita, M.; Nakajima, K.; Ikeya, Y.; Mitsunashi, H. Polyacetylenes from the roots of *Panax ginseng*. *Phytochemistry* **1991**, *30*, 3327–3333.
 - (31) Hirakura, K.; Morita, M.; Nitsu, K.; Ikeya, Y.; Maruno, M. Linoleoylated polyacetylenes from the roots of *Panax ginseng*. *Phytochemistry* **1994**, *35*, 963–967.
 - (32) Baranska, M.; Schulz, H.; Christensen, L. P. Structural changes of polyacetylenes in American ginseng root can be observed in situ by using Raman spectroscopy. *J. Agric. Food Chem.* **2006**, *54*, 3629–3635.
 - (33) Lam, J.; Christensen, L. P.; Færch, T.; Thomsen, T. Acetylenes from *Solidago* species. *Phytochemistry* **1992**, *31*, 4159–4161.
 - (34) Christensen, L. P.; Kreutzmann, S. Determination of polyacetylenes in carrot roots (*Daucus carota* L.) by high-performance liquid chromatography coupled with diode array detection. *J. Sep. Sci.* **2007**, *30*, 483–490.
 - (35) Purup, S.; Larsen, L. B. Integrated proteomics and biological assays for characterization of non-nutritional effects of butyric acid on intestinal cells. *J. Anim. Feed Sci.* **2007**, *16*, 163–171.
 - (36) Purup, S.; Vestergaard, M.; Pedersen, L. O.; Sejrsen, K. Biological activity of bovine milk on proliferation of human intestinal cells. *J. Dairy Res.* **2007**, *74*, 58–65.

Received May 9, 2009. Revised manuscript received August 3, 2009. Accepted August 4, 2009. This research was supported by the EU-Interreg III A programme in the Fyn–KERN Region, the Danish Council for Strategic Research (project No. 2101-01-0065), and the Danish Agency for Science, Technology and Innovation.